

WON'T HARM
LAWNS*

*when used as directed

IMAGE[®]
HERBICIDES

CRABGRASS, NUTSEGE[†] & WEED KILLER

FOR LAWNS

READY-TO-SPRAY



Nutsedge



Crabgrass



Dandelion



Clover

**KILLS EVEN TOUGH
WEEDS, ROOTS & ALL!**

KEEP OUT OF REACH OF CHILDREN

CAUTION SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY INFORMATION,
FIRST AID AND COMPLETE DIRECTIONS FOR USE.

Active Ingredients:	By Wt.
Sulfentrazone.....	1.6%
Quinclorac.....	5.0%
Other Ingredients.....	93.4%
	100.0%

This product contains 0.67 lb of active ingredient per gallon of product (0.14 lb ai sulfentrazone and 0.49 lb ai quinclorac)
* Not for use on purple nutsedge in California

300535834 - 08/24 | IM209 V10

TREATS
6,000^{sq.} FT.

NET CONTENTS
24 FL OZ
(710 mL)

IMAGE
HERBICIDES
CRABGRASS, NUTSEDGE[®]
& WEED KILLER
FOR LAWNS
READY-TO-SPRAY

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-0761. To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. If empty: Offer for recycling, if available, or dispose of in trash.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg No. 279-3463-90098
EPA Est. No. 59144-GA-1, 8119-WA-1,
70815-GA-001, 39578-TX-1

Subscript used is first letter of run code on container.

1-800-265-0761
gardendivision@central.com

See inside booklet for additional
Precautionary Information, First Aid and
complete Directions For Use.

Image, Central Garden & Pet and Central
Garden & Pet with design are trademarks of
Central Garden & Pet Company.

©2017-2024 Central Garden & Pet Company

*Not for use on purple nutsedge in California

300535835-8/24-IM209 v10

KILLS WEEDS
FAST

IMAGE[®]
HERBICIDES

CRABGRASS, NUTSEGE[†] & WEED KILLER

FOR LAWN

READY-TO-SPRAY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

APPLICATION INSTRUCTIONS AND PRECAUTIONS

WHERE TO USE 	Lawns: Well established turfgrasses. Cool Season Grasses: Fine & Tall Fescue, Perennial Ryegrass, Kentucky & Rough Bluegrass, Warm Season Grasses: Common & Hybrid Bermudagrass, Buffalo grass, Centipede grass, Seashore Paspalum, Zoysiagrass. Do not use on St. Augustine grass.
KILLS 	Sedges, Grasses and Broadleaf Weeds: Yellow Nutsedge, Kyllinga, Crabgrass, Foxtail, Ground Ivy, Wild Violet, Plantain, Dandelion, Clover, Dollarweed, Spurge, Woodsorrel, Knotweed, Chickweed, Curly Dock, Wild Onion, Wild Garlic, and other listed weeds.
WHEN TO USE	When weeds are small and actively growing. For best performance, apply when temperatures are between 55°F - 85°F.
HOW MUCH TO USE	4 fl.oz. per 1 gallon of water per 1,000 ft ² . Spot spray weed directly until thoroughly covered. Mixes automatically as you spray. Some weeds may require repeat application. (See detailed instructions.)
COVERAGE	24 fl.oz. covers up to 6,000 sq.ft.

See inside booklet for additional Precautionary Information, First Aid and complete Directions For Use.



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 279-3463-90098
EPA Est. No. 59144-GA-1, 819-WA-1,
70815-GA-001, 39578-TX-1

Subscript used is first letter of run code
on container.

1-800-265-0761
gardendivision@central.com

*Not for use on purple nutsedge in California

Image, Central Garden & Pet, and Central Garden & Pet with
design are trademarks of Central Garden & Pet Company.
© 2015-2024 Central Garden & Pet Company

300535835-8/24-IM209 v10



8 13576 00690 7

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants in adjacent areas. Do not contaminate water when disposing of equipment washwaters or rinsates.

Groundwater advisory: The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soil classified as sand which have less than 1% organic matter.

Surface water advisory: This herbicide can contaminate surface water through spray drift. Under some conditions, this product may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

PRODUCT INFORMATION

This product selectively controls tough weeds like crabgrass, chickweed, dandelion, clover, ground ivy, purple t and yellow nutsedge (nutgrass), Kylinga, wild onion and garlic, broadleaf plantain, purslane, wild violet, spurge, and other listed annual, biennial, and broadleaf weeds in a variety of lawns, including:

* Not for use in California

Cool Season Grasses
Tall & Fine Fescue, Kentucky Bluegrass, Rough Bluegrass, Ryegrass (perennial)
Warm Season Grasses
Common & Hybrid Bermudagrass, Zoysiagrass, Centipedegrass, Buffalograss, Seashore Paspalum

This product has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties have been evaluated. Some turf varieties are more susceptible to temporary turf discoloration than others. It is recommended to treat a small area prior to treatment of larger areas.

This unique formula causes visual symptoms to develop quickly within 3 to 7 days after application. This product involves uptake by both weed roots and shoots. Complete kill should

occur in 2-3 weeks. This product also controls yellow nutsedge tubers (nutlets) in the soil over time. A postemergence application of this product is improved when adequate soil moisture is present at application.

HOW TO APPLY:

Ready to Spray Hose-End Sprayer

1. Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
2. See below for directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the entire measured area.

Hose End Sprayer Connection

1. Shake well.
2. Make sure control lever is in the **OFF** position. Attach to hose.
3. Turn on water.

Spray

1. Aim towards direction of application and away from you.
2. Turn control lever to **ON** to begin spraying.
3. Spray evenly and as directed for specified type of treatment area.

Stop

1. When finished turn control lever to **OFF**
2. Turn faucet **OFF**. Discharge residual water pressure by turning control lever **ON** and **OFF** again.
3. Disconnect sprayer and store / discard as instructed.

COVERAGE- 4 fl.oz. per 1000 sq.ft.

APPLICATION INSTRUCTIONS:

- Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application.
- Listed sedges and grassy weeds: Spray directly on emerged weeds. This product is effective against newly emerged and established sedges and grassy weeds.
- Listed broadleaf weeds: Spray directly on newly emerged weeds.
- For best results, apply to weeds when less than 3 inches high.
- Apply in calm weather.
- Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces.
- Good spray coverage is needed for optimum control.
- Avoid overlap spraying.
- Avoid side to side or swinging arm motion, as it can result in uneven coverage.
- If listed weeds are not controlled by the initial application, apply a second application 4 to 6 weeks after the initial treatment.
- Water lawn with at least 0.5 inches within 7 days (but not sooner than 1 day after application) if no rainfall or irrigation occurs within 7 days after application.

APPLICATION RESTRICTIONS

Do not exceed 5.1 fl oz per 1000 ft² per application.

Keep children and pets out of the treated area until sprays have dried.

Do not spray over the top of flowers, ornamentals (including herbaceous) or garden vegetables.

Do not use on turfgrasses other than those listed on the label.

Do not use in greenhouses.

Do not graze or feed animals with treated grass clippings.

Do not apply to newly planted, plugged, or sodded turfgrass.

Do not release spray at a height greater than 30 inches above the ground.

Do not apply when wind speeds are greater than 10 mph at the application site

APPLICATION PRECAUTIONS

Use only on well established non-stressed turfgrass. Applications to stressed plants (turfgrass or target weeds) may result in temporary discoloration or poor weed control. Allow turfgrass to develop good root system before applying.

If discoloration occurs, treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turfgrass growth.

Avoid applying when windy to prevent product drifting to other plants.

Avoid reseeding, winter overseeding, or plugging lawns for at least 4 weeks after treatment.

Grass and sedge weed species controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	<i>(Echinochloa crus-galli)</i>	Kyllinga, fragrant	<i>(Kyllinga sesquiflorus)</i>
Broadleaf Signalgrass	<i>(Brachiaria platyphylla)</i>	Kyllinga, tufted	<i>(Kyllinga pumila)</i>
Crabgrass, Large	<i>(Digitaria sanguinalis)</i>	Japanese stiltgrass	<i>(Microstegium vimineum)</i>
Crabgrass, Smooth	<i>(Digitaria ischaemum)</i>	Jungle rice	<i>(Echinochloa colona)</i>
Deergrass	<i>(Panicum cladesium)</i>	Nutsedge, purple ¹	<i>(Cyperus rotundus)</i>
Goosegrass	<i>(Eleusine indica)</i>	Nutsedge, yellow	<i>(Cyperus esculentus)</i>
Foxtail, giant	<i>(Setaria faberi)</i>	Sedge, annual	<i>(Cyperus compressus)</i>
Foxtail, green	<i>(Setaria viridis)</i>	Sedge, globe	<i>(Cyperus globulosus)</i>
Foxtail, yellow	<i>(Setaria glauca)</i>	Sedge, cylindrical	<i>(Cyperus retrorsus)</i>
Kyllinga cocks-comb	<i>(Kyllinga squamulata)</i>	Sedge, Surinam	<i>(Cyperus surinamensis)</i>
Kyllinga green	<i>(Kyllinga brevifolia)</i>	Sedge, Texas	<i>(Cyperus polystachyos)</i>
Kyllinga, false green	<i>(Kyllinga gracillima)</i>		

For optimal control of Goosegrass, Jungle rice, or Purple Nutsedge¹, repeat treatment every 3-4 weeks.

¹ Not for use in California

Broadleaf weeds controlled or suppressed by this product:

Common Name	Scientific Name
Bedstraw, catchweed	<i>(Galium aparine)</i>
Beggarticks ¹	<i>(Bidens frondosa)</i>
Beggarweed, Florida	<i>(Desmodium tortuosum)</i>
Bindweed, field	<i>(Convolvulus arvensis)</i>
Bittercress	<i>(Cardamine spp.)</i>
Black medic	<i>(Medicago lupulina)</i>
Buttercups	<i>(Ranunculus spp.)</i>
Butterweed, common (poorjoe)	<i>(Oxalis tenes)</i>
Carolina geranium	<i>(Geranium carolinianum)</i>
Carpetweed	<i>(Mollugo verticillata)</i>
Chickweed, common	<i>(Stellaria media)</i>
Chickweed, mouseear	<i>(Cerastium vulgatum)</i>
Cinquefoil	<i>(Potentilla spp.)</i>
Clover	<i>(Trifolium spp.)</i>
Clover, ayce	<i>(Alysicarpus ovalifolius)</i>
Clover, hop	<i>(Trifolium campestre)</i>
Clover, red	<i>(Trifolium pratense)</i>
Clover, white	<i>(Trifolium repens)</i>
Cudweed	<i>(Gnaphalium spp.)</i>
Daisy, English	<i>(Bellis perennis)</i>
Dandelion	<i>(Taraxacum officinale)</i>
Dandelion, false	<i>(Microseris cuspidata)</i>
Dayflower, common	<i>(Commelina diffusa)</i>
Dock, Curly	<i>(Rumex crispus)</i>
Dollarweed (pennywort)	<i>(Hydrocotyle spp.)</i>
Eclipta	<i>(Eclipta prostrata)</i>
Evening primrose	<i>(Oenothera biennis)</i>
Fiddleneck	<i>(Amsinckia spp.)</i>
Flaree	<i>(Erodium spp.)</i>
Flaree, redstem	<i>(Erodium cicutarium)</i>
Galinisoga	<i>(Galinsoga ciliata)</i>
Goldenrod	<i>(Solidago spp.)</i>
Ground ivy	<i>(Elechinea hederacea)</i>
Groundsel, common	<i>(Senecio vulgaris)</i>
Henbit	<i>(Lamium amplexicaule)</i>
Knawel	<i>(Scleranthus annuus)</i>
Knotweed, prostrate	<i>(Polygonum aviculare)</i>
Kochia	<i>(Kochia scoparia)</i>
Lambsquarters, common	<i>(Chenopodium album)</i>
Lawn burweed (Spurweed)	<i>(Solinia pterosperma)</i>
Lespedeza, common	<i>(Lespedeza striata)</i>
London Rocket	<i>(Sisymbrium irio)</i>
Mallow, common	<i>(Malva neglecta)</i>
Morning glory	<i>(Ipomoea spp.)</i>
Morning glory, entireleaf	<i>(Ipomoea hederacea var. integruscula)</i>
Morning glory, ivyleaf	<i>(Ipomoea hederacea)</i>
Morning glory, ivyleaf	<i>(Ipomoea wrightii)</i>
Morning glory, pitted	<i>(Ipomoea lacunose)</i>
Morning glory, tall	<i>(Ipomoea purpurea)</i>
Parsley piert	<i>(Alchemilla arvensis)</i>

Broadleaf weeds controlled or suppressed by this product (cont'd):

Common Name	Scientific Name
Pigweed, Redroot	<i>(Amaranthus retroflexus)</i>
Pigweed, Smooth	<i>(Amaranthus hybridus)</i>
Pigweed, Tumble	<i>(Amaranthus albus)</i>
Pineapple weed	<i>(Matricaria matricariodes)</i>
Plantain, Broadleaf	<i>(Plantago comanata)</i>
Plantain, Buckhorn	<i>(Plantago lanceolata)</i>
Plantain, Narrowleaf	<i>(Plantago narrowforma)</i>
Puncturevine	<i>(Tribulus terrestris)</i>
Purslane, common	<i>(Portulaca oleracea)</i>
Pusley, Florida	<i>(Richardia scabra)</i>
Redweed	<i>(Melochia corchorifolia)</i>
Shepherd's-Purse	<i>(Capsella bursa-pastoris)</i>
Smartweed, Pennsylvania	<i>(Polygonum pensylvanicum)</i>
Sorrel, Red	<i>(Rumex acetosella)</i>
Speedwell	<i>(Veronica spp.)</i>
Speedwell, slender	<i>(Veronica filiformis)</i>
Speedwell, thymeleaf	<i>(Veronica serpyllifolia)</i>
Spurge, (annuals)	<i>(Euphorbia spp.)</i>
Spurge, prostrate	<i>(Euphorbia humistrata)</i>
Spurge, spotted	<i>(Euphorbia maculata)</i>
Star-of-Bethlehem	<i>(Ornithogalum umbellatum)</i>
Velvetleaf	<i>(Abutilon theophrasti)</i>
Violet, wild	<i>(Viola pratensis)</i>
Violet, Johnny-Jump-Up	<i>(Viola raietensis)</i>
Wild garlic	<i>(Allium vineale)</i>
Wild onion	<i>(Allium canadense)</i>
Woodsorrel, creeping	<i>(Oxalis corniculata)</i>
Woodsorrel, yellow	<i>(Oxalis stricta)</i>

¹ Not for use in California

Certain listed weeds may require more than one application to achieve season-long control.

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing. Do not store below 32°F. Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat.

In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division: 1-800-265-0761.

To confine spill, if liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. If empty, offer for recycling, if available, or dispose of in trash.

If partially filled, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

