

WON'T HARM
LAWNS*

*when used as directed

IMAGE[®]
HERBICIDES

CRABGRASS & WEED KILLER

FOR LAWNS

CONCENTRATE



Crabgrass



Chickweed



Dandelion



Clover

**KILLS EVEN THE TOUGH WEEDS,
ROOTS & ALL!**

FOR OUTDOOR RESIDENTIAL USE ONLY
KEEP OUT OF REACH OF CHILDREN
CAUTION STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

See attached booklet for complete Precautionary Statements
and Directions for Use, including Storage and Disposal, and back
panel for First Aid.

300535844-05/24 | IM214 V8

ACTIVE INGREDIENTS:
MCPA, dimethylamine salt 4.78%
Quinclorac 1.56%
Dicamba, dimethylamine salt 0.43%
OTHER INGREDIENTS 93.23%
TOTAL 100.00%

THIS PRODUCT CONTAINS:
0.338 lb 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 3.90%
0.136 lb 3,4-dichloro-4'-quinclinoic acid per gallon or 1.56%
0.031 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.36%

1 QUART TREATS
4,000 SQ. FT.

NET CONTENTS
1 QUART
(32 FL OZ) (946 mL)

IMAGE[®]
HERBICIDES

CRABGRASS & WEED KILLER

FOR LAWNS

CONCENTRATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment wash waters or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



Manufactured for:
Central Garden & Pet, Garden Division
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 2217-954-90098
EPA Est. No. 59144-GA-1, 92162-MD-1,
Subscript used is first letter of run code on container.

1-800-265-0761
gardendivision@central.com
www.imageforweeds.com

Image, Central Garden & Pet and Central
Garden & Pet with design are trademarks of
Central Garden & Pet Company.

© 2017-2024 Central Garden & Pet Company

300535845 - 5/24 | IM214 V8

[illegible]

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Use Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Apply when daytime temperatures are between 50°F and 90°F. Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur. For bermudagrass lawns do not apply when daytime temperatures exceed 85°F.
- Do not apply at rates above those specified on this label.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

SHAKE WELL BEFORE USING

2. How it Works

This product contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

3. Where to Use

This product can be applied to the following established turfgrass species:	
Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass, Ryegrass, perennial and annual, Tall fescue, Bentgrass, Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	Bermudagrass, common and hybrid, Zoysiagrass, Buffalograss
Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.	
DO NOT apply this product to the following:	
Warm Season Turfgrass	Other Areas
St. Augustinegrass, Bahia grass, Centipede grass, Carpet grass, Seashore paspalum	Lawns with desirable clovers or legumes, Dichondra, Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Turfgrass species that are not listed on this label, Gardens including vegetables, fruit trees, vines and berries.

4. When to Use:

Apply this product when weeds are small and actively growing.

Crabgrass (large and smooth):

- Spring: Apply this product from crabgrass emergence until the crabgrass is 3 to 4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at the same rate at 21 days later for dense populations.

Foxtail (green, giant, and yellow):

- Spring and early summer: Apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at the same rate at 21 days later for dense populations.

Broadleaf Weeds:

- Spring: Apply when dandelion, plantain, buttercup, speedwell, knotweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- Fall: Apply when henbit, chickweed, white clover, ground ivy, lawn burweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

5. How Much to Use

Limitations on broadcast applications to residential turfgrass
The maximum application rate is 8 fl.oz. of product per 1,000 sq.ft. per application (0.92 lb MCPA acid equivalent and 0.08 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 16 fl.oz. of product per 1,000 sq.ft. (1.84 lb MCPA ae and 0.17 lb dicamba ae per acre acid equivalent per acre).

5.1 Entire Lawn Treatments

THREE EASY STEPS TO MIX AND APPLY:

- Measure the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- For hose-end sprayers: Set the dial to the RATE PER GALLON and keep the dial set to this setting while spraying. The RATE PER GALLON for this product is 4 fl.oz. of product per gallon. See instructions below.
- For hand-operated sprayers (backpack sprayers, compression (pump-up) sprayers, knapsack sprayers or tank type): The RATE PER 1,000 sq.ft. for this product is 8 fl.oz. of product per 1,000 sq.ft. of lawn. Measure and mix the proper amount of product with water as described below.
- Spray evenly over the lawn.

5.2 Spot Treatments for Small Areas or Individual Weeds

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up): Mix 4 fl.oz. of product with 1 gallon of water. Note: Approximately one (1) gallon of spray solution will treat 500 sq.ft. See below for additional spray preparations. Shake well before spraying. Apply to actively growing weeds during the early growth stages. Spray coverage should be uniform and complete.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Approximate Treated Area, Square Feet
4 fl. oz.	1 gallon	500 sq.ft.
6 fl. oz.	1.5 gallons	750 sq.ft.
8 fl. oz.	2 gallons	1,000 sq.ft.
12 fl. oz.	3 gallons	1,500 sq.ft.
Equal measures: 1 cup = 8 fl. oz. = 0.5 pint 4 fl. oz. = 8 tablespoons		

6. Cultural Tips:

For newly seeded areas:

- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged, or plugged areas:

- The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding:

- Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- Adequate moisture and mild environmental conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) at 24 hours before and after the application.
- Avoid applications of this product when turfgrasses are under stress since injury may result.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

7. Weeds Controlled

GRASS WEEDS	Crabgrass, large & smooth Foxtail, green, yellow & giant	Kikuyugrass Signalgrass, broadleaf
Annual yellow sweetclover Aster Austrian fieldcress Bedstraw Beggarticks Betony, Florida Bindweed, field Bird vetch Bitter wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Bloodflower milkweed Blue lettuce Blue vervain Bracted plantain Brassbuttons Bristly oxtongue Broadleaf dock Broadleaf plantain	Broomweed Buckhorn Buckhorn plantain Bulbous buttercup Bull thistle Bull nettle Burclover Burdock Burning nettle Burweed Buttercup Buttonweed Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catnip Catsear Chickweed, common Chickweed, mouseear Chicory Cinquefoil Clover, crimson Clover, hop	Clover, red Clover, strawberry Clover, sweet Clover, white Cockle Cocklebur Common mullein Creeping Jenny (Creeping Charlie) Cudweed Curly dock Daisy, English Daisy, Reabane Daisy, oxeye Dandelion Dichondra Dogbane Dogfennel Dogweed Elderberry False dandelion False fax False sunflower Fiddleneck Florida pusley

7. Weeds Controlled

Frenchweed	Pepperweed	Strawberry, India mock
Galinsoga	Pigweed	Tall nettle
Goathead	Pineywoods bedstraw	Tall vervain
Goldenrod	Plains coreopsis	Tansy ragwort
Ground Ivy	(tickseed)	Tansy mustard
Gumweed	Plantain	Tanweed
Hairy fleabane	Poison ivy	Thistle
Hawkweed	Poison oak	Trailing crownvetch
Headail	Pokeweed	Tumble mustard
Heartleaf drymary	Poorjoe	Tumble pigweed
Heathaster	Prairie sunflower	Velvetleaf
Hedge bindweed	Prickly lettuce	Venice mallow
Hedge mustard	Prickly sida	Virginia buttonweed
Hemp	Prostrate knotweed	Virginia creeper
Henbit	Prostrate pigweed	Virginia pepperweed
Hoary cress	Prostrate spurge	Wavyleaf bulblet
Hoary plantain	Prostrate vervain	Western clematis
Hoary vervain	Puncturevine	Western salsify
Horsenettle	Purslane, common	White mustard
Jimsonweed	Ragweed	Wild aster
Knawel	Red sorrel	Wild buckwheat
Knotweed	Redroot pigweed	Wild carrot
Kochia	Redstem filaree	Wild four-o'clock
Lambsquarters	Rough cinquefoil	Wild garlic
Lespedeza	Rough fleabane	Wild geranium
Mallow	Russian pigweed	Wild lettuce
Matchweed	Russian thistle	Wild marigold
Mexicanweed	Scarlet pimpernel	Wild mustard
Milk vetch	Scotch thistle	Wild onion
Morningglory	Sheep sorrel	Wild parsnip
Mouseear hawkweed	Shepherdspurse	Wild radish
Mugwort	Slender plantain	Wild rape
Musk thistle	Smallflower galinsoga	Wild strawberry
Mustard	Smooth dock	Wild sweet potato
Narrowleaf plantain	Smooth pigweed	Wild vetch
Narrowleaf vetch	Sorrel	Wild violet
Nettle	Sowthistle	Woodsorrel
Orange hawkweed	Spanishneedles	Woolly croton
Oriental cocklebur	Speedwell	Woolly morningglory
Oxalis	Spiny amaranth	Woolly plantain
Parsley-piert	Spiny cocklebur	Wormseed
Parsnip	Spiny sowthistle	Yarrow
Pearlwort	Spotted catsear	Yellow rocket
Pennycress	Spotted spurge	Yellowflower pepperweed
Pennywort	Spurweed	
Peppergress	Stinging nettle	

NOTES

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable containers: Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Image, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

NOTES

[illegible]

NOTES

[illegible]