



FOR LAWNS

CONCENTRATE









Nutsedge

Crabgrass

Dandelion

Clover

KILLS EVEN TOUGH WEEDS, ROOTS & ALL!

KEEP OUT OF REACH OF CHILDREN CAUTION SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY INFORMATION, FIRST AID AND COMPLETE DIRECTIONS FOR USE.

Active Ingredients:	By Wt.
Su l fentrazone	1.6%
Quinclorac	
Other Ingredients	
	100.00

This product contains 0.57 lb of active ingredient per gallon of product (0.14 lb ai sulfentrazone and 0.43 lb ai quinclorac)
† Not for use on purple nutsedge in California

TREATS 6,000 SQ. **NET CONTENTS** 4 FL OZ (710 mL)



CONCENTRATE

/ FIRST AID				
Ifswallowed	Call poison control center or doctor immediately for treatment advice, Have a person sip a glass of water if able to swallow. Do not induce vemiting unless told to do so by the poison control center or doctor, Do not give anything by mouth to an unconscious person.			
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.			
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.			
	HOTLINE NUMBER			
	ontainer or label with you when calling a poison control center or doctor, or going for r also contact 1-800-265-0761 for emergency medical treatment information.			

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet, Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark, Drift and runoff may be hazardous to plants in adjacent areas, Do not contaminate water when disposing of equipment washwaters or rinsates.

Groundwater advisory: The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of label use, Use of this chemical in areas where soils are permeable, particularly where

the water table is shallow, may results in groundwater contamination. Do not use on coarse soil classified as sand which have less than 1% organic matter.

<u>Surface water advisory</u>: This herbicide can contaminate surface water through spray drift. Under some conditions, this product may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

STORAGE AND DISPOSAL

Pesticide Storage

Keep from freezing, Do not store below 32°F, Keep out of reach of children and animals, Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. In case of a spill, avoid contact, its loade area and keep out animals and unprotected persons. Confine spills, Call Central Barden & Pet, Barden Division: 1-800-265-0761.

Confine spill, If liquid, dike surrounding area or absorb with and, call titer or commercial day, if dry material, cover to prevent dispersal. Place damaged package in a holding container, Identify contents.

PESTICIOE DISPOSAI

Pesticide waste are cutely hazardous.Improper disposal of excess pesticide, spray mixture, or rinsate is a vidation of Federal Law, If these wastes cannot be disposed of by use according to label instructions, contact your State Pestidide or Frommental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling
Non-refillable container. Do not reuse or refill this container. <u>If empty:</u> Offer for recycling, if available, or

In a remaining containing to interest or term and containing <u>framining</u> other to recogning in available, or dispose of in trash.

<u>If partially filled:</u> Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



gardendivision@central.com

Manufactured for: Central Garden & Pet, Garden Division 1000 Parkwood Circle, Suite 700 Atlanta, GA 30339

† Not for use on purple nutsedge in California Image, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

NOTES

_





FOR LAWNS

CONCENTRATE

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling, Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

APPLICATION INSTRUCTIONS AND PRECAUTIONS

WHERE TO USE
M

iwns:

Well established turfgrasses. Cool Season Grasses: Fine & Tall Fescue, Perennial Ryegrass, Kentucky & Rough Bluegrass.

Warm Season Grasses: Common & Hybrid Bermudagrass, Buffalograss, Centipedegrass, Seashore Paspalum, Zoysiagrass. Do not use on St. Augustinegrass.

KILLS

HOW MUCH TO USE Sedges, Grasses and Broadleaf Weeds: Yellow Nutsedge, Kyllinga, Crabgrass, Foxtail, Ground Ivy, Wild Violet, Plantain, Dandelion, Clover, Dollarweed, Spurge, Woodsorrel, Knotweed, Chickweed, Curly Dock, Wild Onion, Wild Garlic, and other listed weeds.

when weeds are small and actively growing. For best performance, apply when temperatures are between 55°F- 85°F.

apply when temperatures are between 55°F-85°F.

4fl.oz.per 1 gallon of water per 1,000 ft². Spot spray weed directly until thoroughly covered.

Some weeds may require repeat application. (See detailed instructions.)

COVERAGE 24 fl.oz. covers up to 6,000 sq.ft.

 $See \ inside \ booklet \ for \ additional \ Precautionary \ Information, First \ Aid \ and \ complete \ Directions \ For \ Use.$



Manufactured for: Central Garden & Pet, Garden Division 1000 Parkwood Circle, Suite 700 Atlanta, GA 30339

EPA Reg No. 279-3463-90098 EPA Est. No. 59144-GA-1_A, 8119-WA-1_B, 70815-GA-001_C 39578-TX-1.

Subscript used is first letter of run code on container.

1-800-265-0761 gardendivision@centra**l.**co † Not for use on purple nutsedge in California Image, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company, © 2015-2024 Central Garden & Pet Company

300535837 - 8/24 | IM210 V9



	FIRST AID
If swallowed	- Cal poison control center or doctor immediately for treatment advice. - Have a person sip a glass of water if able to swalfav. - Do not induce vomiting unless told to do so by the poison control center or o
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
lf in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes, Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
	HOTLINE NUMBER
	t container or label with you when calling a poison control center or doctor, or going for lay also contact 1-800-265-0761 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CALITION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes, socks and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants in adjacent areas. Do not contaminate water when disposing of equipment washwaters or rinsates.

Groundwater advisory: The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may results in groundwater contamination.

Do not use on coarse soil classified as sand which have less than 1% organic matter. Surface water advisory: This herbicide can contaminate surface water through spray drift. Under some conditions, this product may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label. The use of this product not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination. PRODUCT INFORMATION

This product selectively controls tough weeds like crabgrass, chickweed, dandelion, clover, ground ivy, purple[†] and yellow nutsedge (nutgrass), Kyllinga, wild onion and garlic, broadleaf plantain, purslane, wild violet, spurge, and other listed annual, biennial, and broadleaf weeds in a variety of lawns, including:

† Not for use in Ca**l**ifornia

COOL 2692011 GL92262
Tall & Fine Fescue, Kentucky Bluegrass, Rough Bluegrass, Ryegrass (perennial)
Warm Season Grasses
Common & Hybrid Bermudagrass, Zoysiagrass, Centipedegrass, Buffalograss, Seashore Paspalum

This product has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties have been evaluated. Some turf varieties are more susceptible to temporary turf discoloration than others. It is recommended to treat a small area prior to treatment of larger areas. This unique formula causes visual symptoms to develop quickly within 3 to 7 days after application. This product involves uptake by both weed roots and shoots. Complete kill should occur in 2-3 weeks. This product also controls yellow nutsedge tubers (nutlets) in the soil over time. A postemergence application of this product is improved when adequate soil moisture is present at application.

HOW TO APPLY:

- · Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
- Shake well before use.
- Remove measuring cap and fill cap according to chart below.
- Pour into sprayer and add corresponding amount of water.

Amount of Product	Amount of Water	Coverage
1 f l. oz.	1 Quart (32 oz)	250 sq.ft.
2 f l. oz.	1/2 Gallon (64 oz)	500 sq.ft.
4 f l. oz.	1 Ga ll on (128 oz)	1000 sq.ft
6 f l. oz.	1½ Gallons (192 oz)	1500 sq.ft.
8 f l. oz.	2 Ga ll ons (256 oz)	2000 sq.ft.

PUMP-STYLE TANK SPRAYERS

- Close sprayer tightly and pressurize with hand pump.
- · For spot applications, use a coarse spray nozzle. For larger applications, use a fan spray
- Hold tip of wand 12 inches from target area while spraying.

DIAL-STYLE HOSE-END SPRAYERS

- Fill sprayer jar with enough product for the area to be treated, DO NOT ADD WATER,
- Close sprayer, set dial, hook to garden hose and apply.
- Pour any unused product back into the product's original container.

APPLICATION INSTRUCTIONS:

- Best weed control results will be obtained when no rainfall or irrigation occurs within 24 hours after application.
- Listed Sedges and Grassy Weeds: Spray directly on emerged weeds. This product is effective against newly emerged and established sedges and grassy weeds.
- Listed Broadleaf Weeds: Spray directly on newly emerged weeds.
- For best results, apply to weeds when less than 3 inches high.
- Apply in calm weather.
- Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces.
- Good spray coverage is needed for optimum control.
- Avoid overlap spraying.
- Avoid side to side or swinging arm motion, as it can result in uneven coverage.
- If listed weeds are not controlled by the initial application, apply a second application 4 to 6 weeks after the initial treatment.
- · Water lawn with at least 0.5 inches within 7 days (but not sooner than 1 day after application) if no rainfall or irrigation occurs within 7 days after application.

APPLICATION RESTRICTIONS

Do not exceed 5.1 fl oz per 1000 ft² per application.

Keep children and pets out of the treated area until sprays have dried.

Do not spray over the top of flowers, ornamentals (including herbaceous) or garden vegetables.

Do not use on turfgrasses other than those listed on the label.

Do not use in greenhouses.

Do not graze or feed animals with treated grass clippings.

Do not apply to newly planted, plugged, or sodded turfgrass. Do not release spray at a height greater than 30 inches above the ground.

Do not apply when wind speeds are greater than 10 mph at the application site.

APPLICATION PRECAUTIONS

Use only on well-established non-stressed turfgrass. Applications to stressed plants (turforass or target weeds) may result in temporary discoloration or poor weed control. Allow turfgrass to develop good root system before applying.

If discoloration occurs, treated turfgrass will recover with new growth and discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turfgrass growth.

Avoid applying when windy to prevent product drifting to other plants.

Avoid reseeding, winter overseeding or plugging lawns for at least 4 weeks after treatment.

Grass and sedge weed species controlled or suppressed by this product:

Common Name	Scientific Name	Common Name	Scientific Name \
Barnyardgrass	(Echinochloa crus-galli)	Ky l linga,	(Kyllinga gracillima)
Broadleaf	(Brachiaria platyphylla)	false green	
Signa l grass		Kyllinga, fragrant	(Kyllinga sesquiflorus)
Crabgrass, Large	(Digitaria sanguinalis)	Kyllinga, tufted	(Kyllinga pumila)
Crabgrass, Smooth	(Digitaria ischaemum)	Japanese stiltgrass	(Microstegium vimineun
Deergrass	(Panicum cladestinum)	Junglerice	(Echinochloa colona)
Goosegrass	(Eleusine indica)	Nutsedge, purple†	(Cyperus rotundus)
Foxtail, giant	(Setaria faberi)	Nutsedge, yellow	(Cyperus esculentus)
Foxtail, green	(Setaria viridis)	Sedge, annual	(Cyperus compressus)
Foxtail, yellow	(Setaria glauca)	Sedge, globe	(Cyperus globulosus)
Kyllinga,	(Kyllinga squamulata)	Sedge, cylindrical	(Cyperus retrorsus)
cocks-comb		Sedge, Surinam	(Cyperus surinamensis)
Kyllinga, green	(Kyllinga brevifolia)	Sedge, Texas	(Cyperus polystachyos)

For optimal control of Goosegrass, Junglerice, or Purple Nutsedge¹, repeat treatment every 3-4 weeks. † Not for use in California

Common Name	Scientific Name	Common Name	Scientific Name
Bedstraw, catchweed	(Galium aparine)	Morning glory	(Ipomea spp.)
Beggarticks†	(Bidens frondosa)	Morning glory.	(Ipomoea hederacea
Beggarweed, Florida	(Desmodium tortuosum)	entire l eaf	var. integriuscula)
Bindweed, fie l d	(Convolvulus arvensis)	Morning glory, ivyleaf	(Ipomoea hederacea)
Bittercress	(Cardamine spp.)	Morning glory,	(Ipomoea wrightii)
Black medic	(Medicago lupuline)	palmleaf	()
Buttercups	(Ranunculus spp.)	Morning glory, pitted	(Ipomoea lacunose)
Buttonweed, common	(Diodia teres)	Morning glory, tall	(Ipomoea purpurea)
(poorioe)		Pars l ey piert	(Alchemilla arvensis)
Carolina geranium	(Geranium carolinianum)	Pigweed, Redroot	(Amaranthus
Carpetweed	(Mollugo verticillata)	,	retroflexus)
Chickweed, common	(Stellaria media)	Pigweed, Smooth	(Amaranthus
Chickweed, mouseear	(Cerastium vulgatum)		hybridus)
Cinquefoi l	(Potentilla spp.)	Pigweed, Tumble	(Amaranthus albus)
Clover	(Trifolium spp.)	Pineapple weed	(Matricaria
llover, alyce	(Alysicarpus ovalifolius)	l ''	matricariodes)
Jover, hop	(Trifolium campestre)	Plantain, Broadleaf	(Plantago comanalata)
Jover, red	(Trifolium pretense)	Plantain, Buckhorn	(Plantago lanceolata)
Jover, white	(Trifolium repens)	Plantain, Narrowleaf	(Plantago
Cudweed	(Gnaphalium spp.)		narowiforma)
Daisy, Eng l ish	(Bellis perennis)	Puncturevine	(Tribulus terrestris)
Dande l ion	(Taraxacum officianale)	Purslane, common	(Portulaca oleracea)
Dande l ion, fa l se	(Microseris cuspidate)	Pusley, Florida	(Richardia scabra)
Dayf l ower, common	(Commelina diffusa)	Redweed	(Melochia
Oock, Cur l y	(Rumex crispus)		corchorifolia)
Do ll arweed	(Hydrocotyle spp.)	Shepherd's-Purse	(Capsella
(pennywort)			bursa-pastoris)
clipta	(Eclipta prostrata)	Smartweed,	(Polygonum
evening primrose	(Denother biennis)	Pennsy l vania	pensylvanicum)
idd l eneck	(Amsinckia spp.)	Sorrel, Red	(Rumex acetosella)
laree	(Erodium spp.)	Speedwe	(Veronica spp.)
laree, redstem	(Erodium cicutarium)	Speedwe ll , s l ender	(Veronica filiformis)
Ga l insoga	(Galinsoga ciliate)	Speedwell, thymeleaf	(Veronica serpyllifolia)
Goldenrod	(Solidago spp.)	Spurge, (annua l s)	(Euphorbia spp.)
Ground ivy	(Glechmea hederacea)	Spurge, prostrate	(Euphorbia humistrata
Groundse l , common	(Senecio vulgaris)	Spurge, spotted	(Euphorbia maculate)
Henbit	(Lamium amplexicaule)	Star-of-Beth l ehem	(Ornithogalum
Knawel	(Scleranthus annuus)		umbellatum)
(notweed, prostrate	(Polygonum aviculare)	Velvetleaf	(Abutilon theophrasti)
Kochia	(Kockia scoparia)	Violet, wild	(Viola pratincola)
Lambsquarters,	(Chenopodium album)	Violet,	(Viola rafeinesquii)
common	(Calina mtanananana)	Johnny-Jump-Up	(Allium vinanta)
Lawn burweed	(Soliva pterosperma)	Wild garlic	(Allium vineale)
(Spurweed)	(1 d d-/ - d-)	Wild onion	(Allium canadense)
Lespedeza, common	(Lespedeza striafa)	Woodsorrel, creeping	(Oxalis corniculata)
London Rocket	(Sisymbrium irio)	Woodsorrel, yellow	(Oxalis stricta)
Ma ll ow, common	(Malva neglecta)	I	

¹ Not for use in California

Certain listed weeds may require more than one application to achieve season-long control.

STORAGE AND DISPOSAL

Pesticide Storage
Keep from freezing, Do not store below 32°F, Keep out of reach of children and animals,
Store in original container only. Store in a cool, dry place, preferably in a locked storage

area and avoid excess heat.

In case of a spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Central Garden & Pet, Garden Division:

unprotected persons, comine spilist, call central barrein at Pret, Gardein Division: 1-800-265-0761. loconfine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay, If dry material, cover to prevent dispersal. Place damaged package in a holding container, Identify contents.

Pesticide Disposal

Pesticide Uisposal
Pesticide wastes are acutely abzardous. Improper disposal of excess pesticide, spray mixture, or
rinsate is a violation of Federal Law, It these wastes cannot be disposed of by use according to label
instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste
representative at the nearest EPA Regional Office for guidance.

Container Handling
Non-refillable container. Do not reuse or refill this container. If empty: Offer for recycling, if available, or dispose of in trash. If partially filed: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with label directions,

NOTES

_			
_			
-			
-			

NOTES			NOTES	
		-		-
				_
				-
				-
				-
				_
				-
				-
	_			_
				_
				-
	_			-
				_
				-
				=
	_			_
				_
				_
				-
				-
				_